

Volume 13

Number 1

JUNE 2011



# IFE JOURNAL OF SCIENCE

A Journal of the Faculty of Science  
Obafemi Awolowo University, Ile-Ife, Nigeria

ISSN 0794-4896

# Ife Journal of Science

## Aims and Scope

Ife Journal of Science (IJS) aims to publish articles resulting from original research in the broad areas of chemical, biological, mathematical and physical sciences. This extends naturally into frontiers that include the applied areas of Biochemistry and Geology as well as Microbiology and such allied fields as Biotechnology, Genetics, Food Chemistry, Agriculture, Medical and Pharmaceutical Sciences. Shorter-length manuscripts may be accepted as *Research notes*. Review articles on research topics and books are also welcome.

**Editor-in-Chief (Biological Sciences):** Prof. J. O. Faluyi

**Editor-in-Chief (Physical Sciences):** Prof. M. O. Olorunfemi

## Associate Editors

Dr. A. O. Shittu - *Microbiology*  
Prof. G. A. O. Arawomo - *Zoology*  
Prof. A. P. Akinola - *Mathematics*  
Prof. F. O. I. Asubiojo - *Chemistry*  
Prof. A. A. Okunade - *Physics*  
Dr. A. I. Odiwe - *Botany*  
Prof. S. O. Asaolu - *Zoology*  
Prof. O. O. Jegede - *Physics*  
Dr. B. O. Omafuvbe - *Microbiology*  
Dr. F. K. Agboola - *Biochemistry*

Prof. T. O. Obilade - *Mathematics*  
Prof. J. O. Ajayi - *Geology*  
Prof. H. C. Illoh - *Botany*  
Prof. H. B. Olaniyi - *Physics*  
Prof. O. O. Oyedapo - *Biochemistry*  
Prof. S. B. Ojo - *Geophysics*  
Prof. J. O. Nwachukwu - *Geology*  
Dr. A. O. Ogunfowokan - *Chemistry*  
Prof. G. A. Oshinkolu - *CERD*  
Dr. C. C. Adeyemi - *Natural History Museum*

## International Advisory Committee

Prof. Dr. Thomas Foken, *Bayreuth, Germany*  
Prof. Dr. Stefan Wohnlich, *Germany*  
Prof. O. O. Kassim, *Washington DC, USA*  
Dr. Walter Kpikpi, *Tamale, Ghana*  
Prof. J. A. Lockwood, *Laramie, USA*  
Prof. Bjorn Malmgren, *Goteborg, Sweden*  
Prof. Dr. Gunther Matheis, *Berlin, Germany*

Prof. J. O. Nriagu, *Michigan, USA*  
Prof. Kwabena Oduro, *Legon, Ghana*  
Prof. J. O. Olowolafe, *Delaware, USA*  
Prof. Adrian Raftery, *Seattle, USA*  
Prof. L. G. Ross, *Stirling, UK*  
Prof. Reuben H. Simoyi, *Portland, USA*  
Prof. Tetsumaru Itaya, *Japan*

**Two issues of the journal will be published yearly (June and December).**

### **Submission of manuscripts:**

All manuscripts should be submitted to either of the Editors-in-chief, Ife Journal of Science: Prof. J. O. Faluyi, Department of Botany, or Prof. M. O. Olorunfemi, Department of Geology, Obafemi Awolowo University, Ile-Ife, Nigeria.

E-mail: jfaluyi@gmail.com (Tel.: +234-803-7250857)

molorunfel@yahoo.co.uk (Tel.: +234-803-7192169)

## THE EFFECT OF COPPER ON SOME LABORATORY INDICES OF *CLARIAS GARIEPINUS* (BURCHELL 1822).

Obuotor E.<sup>1</sup>; Adewumi A. A.<sup>2\*</sup> and Olaleye V. F.<sup>3</sup>

<sup>1</sup>Dept. of Biochemistry, Obafemi Awolowo University, Ile Ife

<sup>2</sup>Dept. of Biol./Chemical Sciences, University of Sci. & Technology, Ifaki-Ekiti

<sup>3</sup>Dept. of Zoology, Obafemi Awolowo University, Ile Ife

(Received: March, 2011; Accepted: June, 2011)

### ABSTRACT

Haematological studies were carried out on the blood of *Clarias gariepinus* broodstock fed different diets. Five isonitrogenous diets, prepared from locally available feedstuffs were fed to 150 broodfish for 10 weeks in triplicate laboratory aquaria tanks at 10 fish per tank. Four of the diets contain different inclusion levels of copper sulphate tagged CSD1 (1.0mg CuSO<sub>4</sub>/g), CSD2 (2.5mg CuSO<sub>4</sub>/g), CSD3 (5.0mg CuSO<sub>4</sub>/g) and CSD4 (7.5 mg CuSO<sub>4</sub>/g) while the control diet, CSD0 contained no copper sulphate. The haematological parameters measured; haematocrit (HCT), red blood cells, haemoglobin and glucose concentrations were significantly higher ( $p < 0.05$ ) in the broodfish fed CSD0 and CSD1 diets than the other diets. Exposure of *Clarias gariepinus* fish to copper in water, at concentrations above 1.0mg CuSO<sub>4</sub>/g elicits adverse haematological responses and causes homeostatic imbalance. To prevent pollution, it is expedient that there be regulation of the amount of copper sulphate used in cocoa farms, as occurs in south western Nigeria.

**Key words:** *Clarias gariepinus*, Copper sulphate, Haematological, Fish

### INTRODUCTION

In recent years, haematological variables were extensively used when clinical diagnosis of fish physiology was applied to determine the effects of external stressors and disease conditions in fish (Hodson *et al.*, 1978; Dhillion and Gupta, 1983; Benerjec and Kamar, 1988). This, according to Gill and Plant (1981), is as a result of their relationship with energetic (metabolite levels), respiration mechanics (haematological levels) and defence mechanisms (leucocyte levels). Haematological parameters provide an integrated measure of the health status of fishes.

The introduction of toxicants into an environment, where fishes are found, stuns them and/or acts as a stressor of the fish and organisms found in such environment (Olatayo, 2008). The introduction of a toxicant to an aquatic system might decrease the dissolved oxygen concentration, which will impair respiration thus leading to asphyxiation (Warren, 1977). Stickney (1977) also reported that insufficient amount of dissolved oxygen is one of the contributory factors of mortality in some fish species. The darkening of the fish skin, respiratory distress and erratic swimming can be observed when fishes are exposed to acute concentrations of toxicants (De Silva and Ranasinghe, 1989; Ayuba and Ofojekwu,

2002).

Copper sulphate (CuSO<sub>4</sub>) is a popular algicide/fungicide used in cocoa farms. With the recent growing interest in fish farming, a number of earthen ponds are springing up near cocoa farms in the South Western Nigeria. In situations where there are such intercropping with fish farming, unregulated use of copper sulphate can cause leaching into the pond from surrounding farm, which poses toxicological risk to farmed fish and could even lead to massive fish mortality (El-Domiatry, 1987).

Haematological changes shown by fishes exposed to copper include; increase in ammonia (NH<sub>3</sub>) levels, antibody and haematocrit values, haemoglobin level and glucose concentration. Sometimes, the changes are permanent while in other times they are temporary (Olurankinse, 2002). Changes in the haemoglobin concentration of the fish, due to the ingestion of copper may obstruct the uptake of oxygen, thus leading to asphyxiation and eventual death of the fish.

This study was carried out to determine the haematological responses of *Clarias gariepinus* broodstock fed with diets containing different concentrations of copper sulphate, in laboratory aquarium tanks for 10 weeks.

## MATERIALS AND METHOD

The feed samples were prepared with locally available feedstuffs and different levels of copper sulphate (CuSO<sub>4</sub>) incorporations. The trial diets were tagged CSD1 (1.0mg CuSO<sub>4</sub>/g), CSD2 (2.5 mg CuSO<sub>4</sub>/g), CSD3 (5 mg CuSO<sub>4</sub>/g) and CSD4 (7.5 mg CuSO<sub>4</sub>/g) while the control diet, CSD0 has no copper sulphate inclusion (Table 1).

One hundred and fifty male and female *Clarias gariepinus* broodstock were collected from NIFAGOL farms, Yakoyo, Oyo State, and transported to the laboratory in Obafemi Awolowo University, Ile Ife, Nigeria. The fishes

were acclimatized and were all fed the control diet for a period of two weeks after which they were randomly distributed in groups of ten into a plastic aquaria tank (60cmx30cmx30cm) in the laboratory. Each of the trial diets was fed to triplicate groups of the fish for a period of 10 weeks. The physico-chemical parameters of the test media, including the control experiment, were monitored before and during the test period. For the purpose of collecting the blood samples to be used for the haematological studies, an anti-coagulant, ethylene diamine tetra-acetic acid (EDTA) was prepared.

**Table 1:** Experimental Diet Formulations (g/100g).

Formulations	CSD0	CSD1	CSD2	CSD3	CSD4
Soybean	34.2	34.2	34.2	34.2	34.2
Fishmeal	3.6	3.6	3.6	3.6	3.6
Brewery waste	37.8	36.8	35.3	32.8	30.3
Yellow maize	22.5	22.5	22.5	22.5	22.5
Mineral./Vit.	1.0	1.0	1.0	1.0	1.0
Veg. oil	1.0	1.0	1.0	1.0	1.0
Copper sulphate	0	1.0	2.5	5.0	7.5
Total	100.0	100.0	100.0	100.0	100.0

Baseline data on the haematological status of the fish was carried out in triplicates before the experimental feeding commenced. At the end of the culture period, the haematological data were also collected in triplicates, on the fish fed the experimental diets. To collect the fish blood, an incision was made at the dorsal part of the body just behind the head region. The fish body was then mopped with tissue paper to prevent haemolysis due to dilution of oozing blood with any other body fluid. Blood (0.5cm<sup>3</sup>) was then collected with a 2cm<sup>3</sup> glass syringe containing EDTA to give 5mg EDTA per cm<sup>3</sup> of blood properly mixed in a glass tube. The haematological parameters determined include haemoglobin, haematocrit (HCT), red blood cells (RBC), and glucose level.

### HCT determination

HCT was determined immediately after blood collection to prevent swelling of the blood cells, which occurs when blood is exposed to the air (Sniezsko, 1960). Extracted blood was released into a micro centrifuge tube (1.5cm<sup>3</sup>), which was placed in a special micro haematocrit centrifuge

operated at a speed of 5000 rev/min for 10 minutes. Centrifuging separates the plasma from the blood cells. After centrifugation, the haematocrit percentage (HCT) was calculated by measuring the depth of the portion occupied by the red blood cells and dividing by the depth of the whole cell volume.

$$\text{HCT} = \frac{(\text{depth of red blood cells portion})}{\text{depth of the whole portion}} \times 100$$

### RBC determination

The RBC was determined with the aid of a haemocytometer viewed under the microscope to facilitate counting of the red blood cells (Blaxhall and Daisley, 1973). The uncoagulated blood sample was mixed with red blood cell diluting fluid, Dacie's fluid in the ratio of 1:50 (Blood: Dacie's fluid) respectively. The contents were mixed in a tube using an electric mixer after which the contents were allowed to settle for 2-3 minutes. The haemocytometer was covered with a glass tube, which was made to fit it. A centrifuge tube was then filled with the settled blood content and then expelled into the haemocytometer from

the 2 sides. The content was then viewed under the high power (40x) objective of the microscope. The

counts from the sides of hemacytometer were totalled separately and averaged.

$$\text{RBC}/\text{mm}^3 = \frac{(\text{averaged cells counted} \times \text{depth factor (mm)} \times \text{dilution factor})}{\text{area covered (mm)}^2}$$

### Haemoglobin Determination

The haemoglobin content was estimated using the cyanomethaemoglobin method (Larson and Sniezsko, 1961). Blood sample (0.01cm<sup>3</sup>) was mixed together with 2.5cm<sup>3</sup> of the Drabkin's Reagent in a test tube and allowed to settle for 10 minutes at room temperature to allow all the haemoglobin to react with the reagent to form cyanomethaemoglobin. The absorbance of the resultant solution was measured at 540nm using Cam spec visible spectrophotometer.

### Glucose Estimation

The blood sample was centrifuged at a speed of 5000 rev/minutes to obtain the serum, which was used for the estimation of the glucose content. The serum sample (20µl) was added to 2000µl glucose reagent in a test tube. The content was mixed and incubated for 10 minutes at 37°C. A quantity (20µl) of glucose standard solution was also mixed with 200µl of glucose reagent and incubated for 10 minutes at 37°C. The absorbance of the standard solution and those of the samples were measured against those of the reagent blank

in a Cam spec visible spectrophotometer at 546nm (Barham and Trinder, 1972).

## RESULTS AND DISCUSSION

With the increasing concentration of the copper sulphate incorporated into the feed fed to the fish, the total hardness, temperature and pH of the culture medium increased while the dissolved oxygen decreased accordingly (Table 2).

The haematological parameters of the fish fed the control diet were higher at the termination of the experiment when compared to the other treatments. The results of the haematological analysis indicated that increasing concentrations of copper sulphate in the feed of *C. gariepinus* broodstock had pronounced depressive effect on the blood (Table 3). An increase in erythrocyte count was recorded when the fish was fed with diet CSD1 (feed containing 1mg CuSO<sub>4</sub>/g feed). With increasing concentrations of copper in diet of the fish, all the haematological parameters; HCT, glucose, erythrocyte count as well as haemoglobin decreased significantly (p<0.05).

**Table 2:** Mean Values of the Physico-chemical Parameters of the Water in Which *C. gariepinus* Fed the Different Copper Sulphate-incorporated Diets was Raised.

Parameters	CSD0	CSD1	CSD2	CSD3
Temperature (°C)	26.0 ± 0.2 <sup>a</sup>	26.1 ± 0.1 <sup>a</sup>	26.5 ± 0.2 <sup>b</sup>	27.0 ± 0.2 <sup>b</sup>
Dissolved Oxygen (mg/L)	6.0 ± 0.1 <sup>a</sup>	5.8 ± 0.1 <sup>a</sup>	3.2 ± 0.2 <sup>b</sup>	2.1 ± 0.3 <sup>b</sup>
Total hardness (mg/L)	55.4 ± 0.3 <sup>a</sup>	58.5 ± 0.2 <sup>b</sup>	60.3 ± 0.1 <sup>b</sup>	60.5 ± 0.3 <sup>b</sup>
pH	6.6 ± 0.1 <sup>a</sup>	6.7 ± 0.3 <sup>b</sup>	6.8 ± 0.1 <sup>b</sup>	6.8 ± 0.3 <sup>b</sup>

❖ Values with the same superscript in each row are not significantly different from each other (p>0.05)

**Table 3:** Haematological Analysis of the Blood of Fish Fed the Different Copper Sulphate- Incorporated Diets

Parameters	CSD0	CSD1	CSD2	CSD3	CSD4
Haematocrit/HCT	34.1 ± 3.0 <sup>a</sup>	28.3 ± 4.0 <sup>b</sup>	29.1 ± 3.2 <sup>b</sup>	20.5 ± 1.4 <sup>c</sup>	17.7 ± 3.9 <sup>c</sup>
RBC (10 <sup>6</sup> /mm <sup>3</sup> )	2.2 ± 0.5 <sup>a</sup>	2.5 ± 0.2 <sup>a</sup>	1.9 ± 0.6 <sup>b</sup>	2.1 ± 0.3 <sup>b</sup>	1.5 ± 0.1 <sup>c</sup>
Haemoglobin (mmol/L)	8.0 ± 0.6 <sup>a</sup>	6.8 ± 1.1 <sup>b</sup>	6.8 ± 1.3 <sup>b</sup>	6.3 ± 0.2 <sup>c</sup>	5.9 ± 0.3 <sup>c</sup>
Glucose	7.2 ± 1.0 <sup>a</sup>	6.8 ± 1.4 <sup>b</sup>	6.5 ± 2.4 <sup>b</sup>	5.8 ± 0.9 <sup>c</sup>	5.3 ± 0.7 <sup>c</sup>

❖ Values with the same superscript in each row are not significantly different from each other (p>0.05)

The presence of the copper sulphate as a toxicant in this experiment must have caused the increase in total hardness of the water, thus causing the water to retain heat from the atmosphere, leading to increase in the water temperature (Olatayo, 2008). Water temperature and dissolved oxygen are inversely related. Robert (1978) reported that dissolved oxygen generally decreases in solubility with increase in temperature, which also affects other parameters of the aquatic environment that are important to fish health.

As observed in this experiment, high concentration of heavy metals or long-term exposure of fish to sub-lethal concentrations of heavy metals has been shown to decrease various haematological indices and this could result in an anaemic condition of the fish as a result of plasma volume expansion (Kaori *et al.*, 2001). Christenson *et al.* (1972) observed that lower concentrations of copper invariably increase the concentration of haemoglobin in the blood. Increasing levels of copper in the diet is a stressor which will likely cause an osmotic imbalance and change in the fish regulatory system of ionic interchange and which ultimately diminishes the blood pH and decrease the volume of erythrocytes and subsequently lower the percent of haematocrit (Vosyliene, 1999).

The decrease in the haemoglobin of the blood of the fish exposed to increasing concentrations copper in diet, would most probably affect the oxygen uptake and transport. Reduction in the red blood cell count with increasing levels of copper in diet of the fish will cause haemolytic or erythropoietic anaemia in the fish leading to rheological attractions (Kaori *et al.* 2001). Glucose has been shown to be one of the most sensitive indices of the stress state of fish (Vosyliene, 1996; Singh *et al.* 2008). A high concentration of glucose in the blood of the fish fed the control diet, CSD0, is an indication that the fish is not stressed while the decreasing level in the blood of fish fed increasing concentration of  $\text{CuSO}_4$  is a physiological stress indicator.

A decreasing concentration of blood glucose indicates an exhaustion of energy (glycogen) resources and subsequently, the worsening of the status of the fish condition. This result corroborates Vosyliene (1996) who also observed that during long-term tests, high concentration of copper caused a decrease in the concentration of glucose. El-Domiatty (1987) also reported that exposure of *Clarias lazera* to copper results in

haemolysis and anaemia. Van Vuren (1994) also reported that the exposure of *C. gariepinus* to different background copper concentrations induced significant decrease in the erythrocyte count as well as haemoglobin concentrations.

## CONCLUSION

From this experiment, it can be concluded that *C. gariepinus* haematological parameters are affected both by exposure to sublethal concentration of copper sulphate and by the duration of exposure. Exposure of the fish to copper in water, at concentrations above 1.0mg/g will elicit adverse haematological responses and can cause reversible and irreversible homeostatic imbalance. To prevent pollution, it is therefore expedient that there be regulation of the amount of copper sulphate used in cocoa farms, as in south western Nigeria, where some fish farms are located.

## REFERENCES

- Ayuba, V.O. and Ofojekwu, P.C. 2002. Acute toxicity of the root extract of Jimson's weed *Datura innoxia* to the African catfish, *C. gariepinus* fingerlings. *J. Aquac. Sci.* 17(2):131-133.
- Barham, D. and Trinder, P. 1972. An improved colour reagent for the determination of blood glucose by the oxidase system. *Analyst* 97:142-146.
- Blaxhall, P. C. and Daisley K. W. 1973. Routine haematological methods for use with fish blood. *Journal of Fish Biology.* p771 - 781.
- Benerjec, V. and Kamar K. 1988. Effect of zinc, mercury and cadmium on erythrocyte and related parameters in the fish *Anabas tentudinius*. *Envir. Ecol.* 3:739-781.
- Christenson, R.H.; Dent G.A. and Tuszynski A. 1972. Valors de referencia: bibliografia 1 9 7 2 - 2 0 0 2 . <http://www.acclc.cat/invitroveritas/vol4/art43.pdf> Retrieved 15<sup>th</sup> of March, 2010.
- De Silva, C.D. and Ranasinghe J. 1989. Toxicity of four commonly used agrochemicals on *Oreochromis niloticus* (L) fry. *Asian Fish. Sci.* 2:135-145.
- Dhillion, S. S. and Gupta A. K. 1983. A clinical approach to study of pollution intoxication in freshwater teleost (*Clarias berachus*). *Water, Air and Soil Pollut.* 20(1):61-63.
- El-Domiatty, N.A. 1987. Stress response of juvenile *Clarias lazera* elicited by Copper. *Comp. Bioch. and Phys.* 88(2): 259-262.

- Gill A. and Plant J. 1981. Effects of fire on plants and animals. <http://learnline.cdu.edu.au/units/sbi263/ecology/population.html> Retrieved 11th March, 2010.
- Hodson P.V., Blunt B.R. and Spry D.T. 1978. Chronic toxicity of water-borne lead and dietary lead to Rainbow trout, *Salmo gairdneri* in Lake Ontario. *Wat. Res.* 12:869-878.
- Kaori, S., Lewart G.A. and Smith T.M. 2001. Research communication. Blood chemistry values of juvenile Red Pacu (*Piaractus brachypomus*). *Vet. Clin. Path.* 30:50-52.
- Larson, A. and Sniezsko S.F. 1961. Comparison of various methods of haemoglobin in trout blood. *Prog. Fish Cult.* 23:8-17.
- Olatayo, V.S. 2008. Haematological responses of *Oreochromis niloticus* (Trevawas) exposed to various concentrations of poisonous extract from two local plants. *Nig. J. Fisheries*, 5(1):23-30.
- Olurankinse, O.O. 2002. Effects of sub-lethal concentrations of copper sulphate on the body and blood composition of the African catfish, *C. gariepinus* (Burchell 1822) brood in laboratory. B.Sc Thesis, Obafemi Awolowo University. p4-6.
- Robert, R.J. 1978. *Fish Pathology*. Cassel Ltd., London. pp1-77.
- Singh, D., Nath K., Trivedi, S.P. and Sharma Y.K. 2008. Impact of copper on haematological profile of freshwater fish, *Channa punctatus*. *J. Env. Biol.* 29(2):253-7
- Sniezsko, S.F. 1960. *Microhaematocrit as a Tool in Fishery Research and Management*. Spec. Sci. Rep. U.S. Fish Wildlife Service. No 341.
- Stickney, R.R. 1977. *Principles of Warm Water Aquaculture*. John Willey and Sons Incorp. U.K. 375p.
- Van Vuren, J.H.J., Van Der Merwe, M. and Du Preez H.H. 1994. The effect of copper on the blood chemistry of *C. gariepinus* (Clariidae). *Ecotox. Envir. Safety*. 29:187-199.
- Vosyliene, M.Z. 1996. Haematological parameters of Rainbow Trout (*O. mykiss*) during short-term exposure to copper. *Ekologia*. 3:12-18.
- Vosyliene, M.Z. 1999. The effects of heavy metals on haematological indices of fish. *Acta Zoologica Lituanica. Hydrobiologia*. 9(2):76-82
- Warren, C.E. 1977. *Biology and Water Pollution Control*. W.B. Sabders and Company, Philadelphia, USA. 434p.

**Ife Journal of Science**  
**Table of Contents: June Edition 2011; Vol. 13, No. 1**

Omotoye Olorode, Sekinat O. Hassan, Olajumoke A. Olabinjo and Idris O. Raimi	Tithonia (Asteraceae) in Nigeria	1
Obuotor E.; Adewumi A. A. and Olaleye V. F.	The Effect of Copper on Some Laboratory Indices of <i>Clarias Gariepinus</i> (Burchell 1822).	11
Salami, B. M. Conte, R. A. and Falebita, D. E.	Geoelectric Evaluation of the Groundwater Potential of Parts of Osogbo, Southwestern, Nigeria	17
Ogunfowokan A.O., Akanni M.S., Ajibola R.O and Ayinde F.O.	Trophic Status and Physico-Chemical Parameters of Three Reservoirs in Osun State Nigeria	27
Oláýíwolá M.A <sup>1</sup> and Odébòdé M.O.	Foraminiferal Distribution of Southwestern Nigeria's Offshore Littoral Sediments: Benthic Faunal Diversity Indices and Patterns	45
Chinedu S.N., Okochi V. I. and Omidiji O.	Cellulase Production by Wild Strains of <i>Aspergillus Niger</i> , <i>Penicillium Chrysogenum</i> and <i>Trichoderma Harzianum</i> Grown on Waste Cellulosic Materials.	57
Bayode S. and Akpoarebe O.	An Integrated Geophysical Investigation of a Spring in Ibuji, Igbara-Oke, Southwestern Nigeria.	63
M. O. Adepoju and J. A. Adekoya	Reconnaissance Geochemical Study of a Part of Igarra Schist Belt, Southwestern Nigeria	75
Adesina, G.O., Akinyemiju, O.A. and Muoghalu, J.I.	Checklist of the Aquatic Macrophytes of Jebba Lake, Nigeria	93
Fasasi, K. A., Malaka, S. L. O. and Amund, O. O.	Studies on the Life Cycle and Morphometrics of Honeybees, <i>Apis Mellifera Adansonii</i> (Hymenoptera: Apidae) In A Mangrove Area of Lagos, Nigeria.	103
A.O. Olorunfemi, K.S. Salahudeen and T.A. Adesiyani	Ground Water Quality in Ejigbo Town and Environs, Southwestern Nigeria	111
Govardhan Singh, R.S; Ogunsina, B.S. and Radha, C.	Protein Extractability from Defatted <i>Moringa Oleifera</i> Lam. Seeds Flour	121
A. M. A. Sakpere	Identification of ISSR Primers for Genetic Analysis of <i>Telfairia Occidentalis</i> Hook F.	129
O. K. Owoade, F. S. Olise, H. B. Olaniyi, I. B. Obioh and E. Bolzacchini	Mass and Energy Audit in a Nigerian Iron and Steel Smelting Factory: An Operational and Efficiency Study.	133
F. A. Oloyede, B. Aponjolosun & A. A. Ogunwole	Reproductive Potentials of a Tropical Fern <i>Cyclosorus Afer</i> (Christ.) Ching (Thelypteridaceae: Pteridopyhte) at Obafemi Awolowo University, Ile Ife, Nigeria	143
M.O.Olawole, L. Msimanga, S.A.Adegboyega & F.A. Adesina	Monitoring and Assessing Urban Encroachment into Agricultural Land - A Remote Sensing and GIS Based Study of Harare, Zimbabwe	149
Benjamin, U.K and Nwachukwu, J.I	Model Compaction Equation for Hydrostatic Sandstones of the Niger Delta. 161	
J.O. Ojo and C.E. Adeeyinwo	Dependence of Vanadium Recovery on Oxidation State in its Solvent Extraction from Hydrochloric Acid Solutions With TRI N Butyl Phosphate	175
Akintorinwa, O. J., Ojo J. S. and Olorunfemi M. O.	Appraisal of the Causes of Pavement Failure along the Ilesa - Akure Highway, Southwestern Nigeria Using Remotely Sensed and Geotechnical Data	185
O. J. Matthew and O. O. Jegede	Modelling Soil Surface Temperature and Heat Flux Using Force-Restore Method at an Agricultural Site in Ile-Ife, Nigeria.	199
Ojo J.F.	On the Theory of One Dimensional Integrated Autoregressive Bilinear Time Series Modelling	209